

of that Aspidium, & found only one showing this anomaly. & that in 2 only of the funds. It is, so far as I have noticed, confined to the fertile funds. The branching is a leafy bifurcation at tip, for not more than an inch or two at tip. I have also another Aspidium, somewhat collected last summer, in which the stipe branches into two funds. If you desire it, I will send you a specimen of the A. acutæ so branched.

Yours very respy.

J. W. Clinton.

Prof. Asa Gray.

Dear Sir:

Buffalo, Oct. 15, 1862.

Sometime since I sent you seeds of Astragalus Canadensis. & afterwards, I think, the few I could collect of A. Cespitosus. I have not been apprized of their reception; nor have one or more queries I addressed to you been answered. Assuming that they my communications have been received, & that your occupations have prevented any notice of them. I would state some facts & ask for advice in behalf of our Society.

We have collected this season, on the foundation of our local herbarium, more than 600 species excluding the Gramineæ, Cyperaceæ, Gramineæ & Lyngbyaceæ - The season is now substantially over, & we ought to commence

the preparation of my Herbarium. Dr. Tracy promised to send us the plan of his Herbarium, but has omitted to do so. My impression is that the better plan is to affix each specimen to a sheet of stiff paper, to collect the species of each genus in a sheet, paper only labelled - & to put the genera into *pentstelis* ~~accordance~~ labeled with the nat. orders. The use of bound books seems to me objectionable, as it facilitates change, addition, & collection for comparison. You will oblige me very much by your advice, which cannot be too particular.

I have not made any very marvellous discoveries since I last wrote you but the following facts ^{& figures} may be worth writing. I find the old *Carduus Marianus* growing here - a rare escape from cultivation - I cannot reduce it to *Carduus* as now defined, nor to any other genus in your Manual. What is it under the present division?

I find also a *Roseda* with pinnatifid leaves var. ~~pin-~~ - sided pod. The flowers are in conspicuous white racemes. & the odor is like freshly filled cloths. It seems to me that it is not *Whitedea* & that it agrees pretty well with the notes of Persoon's *Eudoridium* -

A small plant - 3-5 in. high & almost invariably with a simple stem - bottom un-
der hair the aspect of *Gratidia hispida*. It
may be it - but is much slender - & confined
to one locality. & from a place I had into
an well-opened cotta (I had lost my
magnifier) I think the inside is bearded.
Shall I send you specimens?

Yesterday I went after the *Bistriatum*
lunarioides var. & found what I suppose
to be the ^a or the *obliquum*. Other
discrepancy (purple leaved) growing with
it. I think I wrote you that
I had ~~not~~ found the *Cespitum* ~~acut-~~
~~iculifer~~, in ~~the~~ hills near Bleau,
slightly branched at top. Yesterday
I examined a hundred or more clumps